

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/114446>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2019 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Publisher's statement:


Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Accepted Manuscript

Title: Does care at a cardiac arrest centre improve outcome after out-of-hospital cardiac arrest? – A systematic review

Authors: J. Yeung, T. Matsuyama, J. Bray, J. Reynolds, M.B. Skrifvars



PII: S0300-9572(19)30029-2
DOI: <https://doi.org/10.1016/j.resuscitation.2019.02.006>
Reference: RESUS 7893

To appear in: *Resuscitation*

Received date: 15 December 2018
Revised date: 8 February 2019
Accepted date: 11 February 2019

Please cite this article as: Yeung J, Matsuyama T, Bray J, Reynolds J, Skrifvars MB, Does care at a cardiac arrest centre improve outcome after out-of-hospital cardiac arrest? – A systematic review, *Resuscitation* (2019), <https://doi.org/10.1016/j.resuscitation.2019.02.006>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Does care at a cardiac arrest centre improve outcome after out-of-hospital cardiac arrest? – A systematic review

J Yeung¹

T Matsuyama²

J Bray³

J Reynolds⁴

M B Skrifvars⁵

¹ Warwick Medical School, University of Warwick, United Kingdom

² Department of Emergency Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan

³ Department of Epidemiology and Preventive Medicine, Monash University, Melbourne Australia.

⁴ Department of Emergency Medicine, Michigan State University, Grand Rapids, Michigan, USA

⁵ Department of Emergency Care and Services, University of Helsinki and Helsinki University Hospital, Finland

on behalf of the Education, Implementation and Team Task Force and Advanced Life Support Task Force of the International Liaison Committee on Resuscitation

AIM

To perform a systematic review to answer 'In adults with attempted resuscitation after non-traumatic cardiac arrest does care at a specialised cardiac arrest centre (CAC) compared to care in a healthcare facility not designated as a specialised cardiac arrest centre improve patient outcomes?'

METHODS

The PRISMA guidelines were followed. We searched bibliographic databases (Embase, MEDLINE and the Cochrane Library (CENTRAL)) from inception to 1st August 2018. Randomised controlled trials (RCTs) and non-randomised studies were eligible for inclusion. Two reviewers independently scrutinized studies for relevance, extracted data and assessed quality of studies. Risk of bias of studies and quality of evidence were assessed using ROBINS-I tool and GRADEpro respectively. Primary

outcomes were survival to 30 days with favourable neurological outcomes and survival to hospital discharge with favourable neurological outcomes. Secondary outcomes were survival to 30 days, survival to hospital discharge and return of spontaneous circulation (ROSC) post-hospital arrival for patients with ongoing resuscitation. This systematic review was registered in PROSPERO (CRD 42018093369)

RESULTS

We included data from 17 observational studies on out-of-hospital cardiac arrest (OHCA) patients in meta-analyses. Overall, the certainty of evidence was very low. Pooling data from only adjusted analyses, care at CAC was not associated with increased likelihood of survival to 30 days with favourable neurological outcome (OR 2.92, 95%CI 0.68 to 12.48) and survival to 30 days (OR 2.14, 95%CI 0.73 to 6.29) compared to care at other hospitals. Whereas patients cared for at CACs had improved survival to hospital discharge with favourable neurological outcomes (OR 2.22, 95%CI 1.74 to 2.84) and survival to hospital discharge (OR 1.85, 95%CI 1.46 to 2.34).

CONCLUSIONS

Very low certainty of evidence suggests that post-cardiac arrest care at CACs is associated with improved outcomes at hospital discharge. There remains a need for high quality data to fully elucidate the impact of CACs.

INTRODUCTION

There is wide variability in survival among hospitals caring for patients after resuscitation from out of hospital cardiac arrest (OHCA). OHCA is common, yet survival outcomes are poor, with substantial regional and international variation.[1-5] Survival from OHCA ranges from 8-16.1%. [6] Measures to maximise favourable neurological outcomes are a research priority for both patients and clinicians.[7] Post-resuscitation care, including percutaneous coronary intervention (PCI) and targeted temperature measurement (TTM), is an important component to achieve good neurological outcome. [8, 9]

In most countries, post resuscitation care is not regionalized to specialised hospitals. [10] There is wide variation among hospitals in the availability and type of post resuscitation care, as well as clinical outcomes. [11, 12] Patients with other time-sensitive emergencies (e.g. trauma, acute myocardial infarction and stroke services) are often triaged to centres which offer speciality services and greater provider experience. [13-16] Such centralised specialist services may improve the provision of targeted post-resuscitation care and offer similar benefits and improve patient outcomes after cardiac arrest.

The International Liaison Committee on Resuscitation (ILCOR) last considered the evidence on this topic in 2015 and concluded that specialist cardiac arrest centres (CACs) may be effective despite a lack of high quality data to support their implementation.[17] Previous observational studies have reported an association between transport to CAC and survival to hospital discharge, but there is inconsistency in the hospital factors that are related to optimal patient outcomes. Whilst most experts agree that a CAC should include a 24-hour cardiac catheterisation laboratory, targeted temperature management, and neurological services that offer electrophysiological modalities for monitoring and prognostication, discrepancies remain in the definition of services that constitute a specialist CAC.[18] The objective of this systematic review was to evaluate outcomes of adults with attempted OHCA resuscitation after care in a specialized cardiac arrest centre compared with care in an institute not designated as a specialized cardiac arrest centre.

METHODS

The protocol for this systematic review was registered with PROSPERO (CRD 42018093369) on 12th April 2018. Reporting of the systematic review was in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. [19] The review was commissioned by ILCOR.

Search strategy and selection criteria

Bibliographic databases (Embase, MEDLINE and the Cochrane Library (CENTRAL)) were searched from inception to February 2018 using a combination of index terms and key words relating to the population, intervention and comparator. The search strategy was developed in conjunction with information specialists at St Michael's Hospital Health Sciences Library (see Supplementary Materials Appendix A for sample search strategy). Reference lists of relevant articles were checked, and clinical trial registers (www.clinicaltrials.gov, www.isrctn.com and <http://www.who.int/ictrp/en/>) were searched to identify on-going trials. (Supplementary Materials Appendix B) Endnote 8 (Thomson Reuters) was used to store records and facilitate screening. The search was repeated on 1st August 2018 to identify any additional studies published during the review process.

Study selection

Studies were eligible for inclusion if they met the following pre-defined criteria:

- 1) Population:** Adults with attempted resuscitation after non-traumatic in-hospital or out-of-hospital cardiac arrest.

- 2) **Intervention and Comparator:** Care at a specialised cardiac arrest centre compared with care in a healthcare facility not designated as a specialised cardiac arrest centre
- 3) **Outcomes:** Primary outcomes were: survival at 30 days with favourable neurological outcome (defined as Cerebral Performance Category (CPC) 1 or 2, modified Rankin Scale (mRS) 0-3); and survival at hospital discharge with favourable neurological outcome (defined as CPC 1 or 2, mRS 0-3).[20, 21] Secondary outcomes were: survival at 30 days; survival at hospital discharge and return of spontaneous circulation (ROSC) after hospital admission.
- 4) **Study Designs:** Randomised controlled trials (RCTs) and non-randomised studies (non-randomised controlled trials, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (e.g., conference abstracts, trial protocols) were excluded. Studies reporting paediatric cardiac arrests (≤ 18 years old) and cardiac arrest secondary to trauma were excluded.
- 5) **Timeframe:** All years and all languages were included as long as there was an English abstract.

Studies were selected by two reviewers (JY/TM) independently by title screening and abstract. Full text of selected studies were retrieved and reviewed by two reviewers (JY/TM) independently. Reasons for exclusion were documented. Reference lists of included studies were also screened.

Definition of Cardiac Arrest Centre

Previous literature has described CACs as institutions that have access to a 24-hour cardiac catheterisation laboratory, targeted temperature management in a critical care facility and prognostication using multimodal approach including delayed clinical examination, neuroelectrophysiological measurements and biomarkers.[22] As there is no widely accepted definition of a cardiac arrest center, we expected variation in description of healthcare institutions and terminology.[18] We accepted 'cardiac arrest centre' or 'regionalized cardiac arrest care' or 'high case volume centres' of similar description in the literature.

Data Extraction and Quality Assessment

A piloted and standardised data extraction form was used to record information on study design, patient population and characteristics, sample size, description of CAC, and outcomes. Two reviewers independently (JY/TM) conducted data extraction (study design, study population, outcome measures and study quality). Any disagreement surrounding the selection of a manuscript or data extraction were resolved either by consensus or arbitration by a third reviewer (MS). Two reviewers (JY/TM) assessed risk of bias of individual studies independently, using the Cochrane Collaboration risk of bias tool[23] for randomised controlled trials and the Risk Of Bias In Non-randomised Studies - of Interventions (ROBINS-I tool)[24] for non-randomised studies. Disagreements were resolved by discussion between the two reviewers.

Data analysis and synthesis

We used meta-analyses to synthesise evidence by outcome, where this approach was not precluded. Generic inverse variance method and random-effects model were used to compute the odds ratio (OR) and 95% confidence interval for dichotomous outcomes.[25] We measured statistical heterogeneity using the Higgins I^2 statistic.[26] We contacted study authors for additional data when it was directly relevant to our outcomes of interest. For observational studies where more than one type of analysis was reported, we preferentially used adjusted or matched case analyses, over unadjusted analyses in meta-analyses. Due to perceived substantial clinical heterogeneity, only effect estimates from adjusted analyses were pooled. In addition, the decision was to pool outcomes only if the effects were in the same direction.

We planned *a priori* subgroup analyses comparing outcomes from patients with shockable and non-shockable initial cardiac rhythm; and direct transfer versus secondary transfer to CACs. In addition, we undertook additional sensitivity analyses to test the robustness of our findings. Firstly, we assessed the impact of our hierarchical approach to selection of analysis type in observational studies by testing the use of other analyses on overall findings. Secondly, we attempted to identify the source of heterogeneity by performing *post-hoc* sensitivity analyses of studies that only included patients with prehospital ROSC and also studies with before-and-after study designs.

The certainty of evidence was assessed using Grading of Recommendations Assessment, Development and Evaluation (GRADE) system in GRADEpro Guideline Development Tool (Evidence Prime, Inc., McMaster University).[27, 28] Review Manager version 5.3 (The Cochrane Collaboration, 2014) was used to produce forest plots.[29]

RESULTS

After removal of duplicates, the literature search yielded 3,065 unique references (3061 from original search and 4 additional studies from updated search). After screening, 22 studies (one randomised controlled trial [30], one prospective study, 4 before and after studies and 16 retrospective analyses) fulfilled eligibility criteria and were included in qualitative synthesis. (Figure 1) Studies were conducted in Australia (n=3), North America (n=8), Europe (n=7) and South East Asia (n=4). The kappa value for identifying studies during initial screening was 0.83. Table 1 contains characteristics of included studies.

Data from 22 OHCA studies were included in the qualitative synthesis, with data from 17 observational studies included in meta-analysis. We described but did not include 5 studies in the meta-analysis: one randomised study since it presented only pilot feasibility data (33 patients, main trial is on-going) [30], three observational studies that only included patients who were discharged alive from hospital and reported long-term outcome data [31-33], and one study which examined the impact of a Post Arrest Consult Team on post cardiac arrest care processes and outcomes.[34]

We found no studies reporting data for in-hospital cardiac arrest patients. One study included in-hospital and out-of-hospital cardiac arrest patients but outcomes were not reported separately.[32]

Risk of bias for individual studies

One randomised controlled study reporting pilot outcomes had moderate overall risk of bias due to lack of treatment allocation. (Table 2a) Twenty-one observational studies had either moderate (n=9) or serious (n=12) overall risk of bias. The most common sources of bias were inadequate adjustment of confounding factors, subject selection (e.g. only including survivors), outcome measurement, and missing data. There was little to no bias from result reporting or the classification of and deviation from the intervention. (Table 2b)

Certainty of evidence across studies

The overall certainty of evidence was very low across all outcomes, primarily due to risk of bias and inconsistency. GRADE summary tables are provided in Appendix C (Supplementary Materials).

Primary outcomes

Survival at 30 days with favourable neurological outcomes (critical)

Survival to 30 days with favourable neurological outcomes were reported by three studies (one retrospective cohort study and one before and after study with adjusted analyses, and one before and after study with unadjusted analyses) recruiting 46,164 patients. [35-37] (Figure 2) In pooled data from two studies (n=45,956) care at CACs was not associated with increased likelihood of a favourable

neurological outcomes at 30 days compared to other hospitals (OR 2.92, 95% CI 0.68 to 12.48). There was moderate to substantial heterogeneity (I^2 73%) and certainty of evidence was very low. One study reporting unadjusted data, also found no association.[37]

Survival at hospital discharge with favourable neurological outcomes (critical)

Survival to hospital discharge with favourable neurologic outcome was reported by six studies. Four studies was included in meta-analysis (one retrospective cohort study and one before and after study with adjusted analyses, and two retrospective cohort studies with unadjusted analyses) recruiting a total of 30,080 patients were included in meta-analysis. [30-33] In pooled data from two studies ($n=3,673$), care at a CAC was associated with increased likelihood of surviving to hospital discharge with favourable neurological outcome compared to other hospitals (OR 2.22 95% CI 1.74 to 2.84). (Figure 3) There was no statistical heterogeneity (I^2 0%) and certainty of evidence was assessed as very low. The direction of effect in two studies reporting unadjusted data favoured care at CACs but was no longer significantly after adjusted analyses. [38, 39] Two studies using an indirect measure (discharge destination) also favoured care at CACs for this outcome. [40, 41]

Secondary outcomes

Survival to 30 days (critical)

Survival to 30 days was reported in six studies (two retrospective cohort studies with adjusted analyses, one before and after study and three retrospective cohort studies with unadjusted analyses) recruiting a total of 45,066 patients. [33, 35, 37, 42-44] In pooled data from two studies ($n=2,693$) care at a CAC was associated with increased likelihood of survival to 30 days post admission (OR 2.14 95% CI 0.73 to 6.29). (Figure 4) There was considerable statistical heterogeneity (I^2 86%) and the certainty of evidence was assessed to be very low. Three of the four studies reporting unadjusted data for this outcome report favourable survival for patients admitted to CACs.

Three additional studies examined long-term survival of cardiac arrest patients by focusing explicitly on the outcomes of patients discharged alive from hospital, these were not included in the meta-analysis. All three studies found that CAC admission was associated with better patient survival. [28-30].

Survival to hospital discharge (critical)

Survival to hospital discharge was reported in ten studies (three retrospective cohort studies and two before and after studies with adjusted analyses, and five retrospective analyses with unadjusted analyses) including 42780 patients. [38-40, 45-50] Data from meta-analysis of 5 studies ($n=11,662$)

found care at a CAC was associated with increased likelihood of survival to hospital discharge compared to other hospitals (OR 1.85 95% CI 1.46 to 2.34). (Figure 5) There was moderate statistical heterogeneity (I^2 69%) and certainty of evidence was assessed to be very low. All five studies reporting unadjusted support the association of the pooled analysis.

Return of spontaneous circulation (ROSC) post hospital admission for patients with ongoing CPR (important)

Data on ROSC post hospital admission for patients with ongoing CPR was obtained from authors of two studies (one retrospective cohort study and one before and after study with unadjusted analyses) comprising which reported post hospital admission ROSC rates in patients with ongoing CPR. [35, 36] (Figure 6) Both studies reported significantly higher ROSC rates in CAC group but data was not pooled based on the lack of adjusted data.

Subgroup Analyses and sensitivity analyses

There were insufficient data to perform the pre-planned pooled subgroup analyses of subjects with shockable vs. non-shockable initial cardiac rhythm, and direct transfer vs. secondary transfer to CAC.

Eight studies [33, 36, 37, 45, 48-51] reported outcomes in CAC versus non-CACs stratified by shockable or non-shockable rhythms. Data were too heterogeneous to allow for meta-analysis (Supplementary Materials Table 3). In patients with shockable rhythm, four studies reported improved outcomes at CACs [36, 48, 49, 51], and three studies reported no difference [37, 45, 50]. In patients with non-shockable initial rhythms, CACs were associated with improved outcomes in one study [51] and no difference in two studies [49, 50].

The majority of the included studies compared OHCA patients who were directly transport to a CAC to those transported to a non-CAC. Only three studies examined outcomes in OHCA patients who were transferred to a CAC from a non-CAC [33, 36, 41]. In these studies, the proportion of patients with secondary transfers were small and, although two studies [33, 41] adjusted for available confounders, there is potential for referral bias (whereby the sickest or those most likely to survive were transferred to CACs). A US study [33] reported no difference in long-term survival between primary and secondary transfers to CACs after adjusting for potential confounders (or in unadjusted 30-day survival data provided by the authors). After conducting numerous analyses to adjust for referral bias, but the study consistently reported increased long-term survival in patients cared for at CACs, with the lowest mortality seen in early transfers. An Australian study [41] reported higher (adjusted) survival in patients who were transferred directly to a CAC compared to secondary

transfers. A small subgroup analysis by Tagami [36] found no difference in survival to 30-days with favourable neurological outcome in patients between direct and secondary transfer to CACs.

Sensitivity analysis replacing adjusted analyses with unadjusted analyses were performed for the outcomes of survival to 30 days with favourable neurological outcomes, survival to hospital discharge with favourable neurological outcomes, survival to 30 days and survival to hospital discharge. These replacements made no difference to our overall findings (data not shown). We performed sensitivity analyses to explore the source of heterogeneity by firstly excluding data from studies that transported only patients with pre-hospital ROSC [38, 40, 45, 46] for two outcomes: Survival to hospital discharge with favourable outcomes (2 studies) and Survival to hospital discharge (4 studies). This did not change our overall findings but did reduce statistical heterogeneity in Survival to hospital discharge from I^2 90% to 52% (data not shown). Secondly, we performed sensitivity analyses by excluding studies of before and after design [36, 37, 48, 49] for three outcomes: Survival to hospital discharge with favourable outcome (1 study), Survival to 30 days (1 study), and Survival to hospital discharge (2 studies). The results were again similar to our main analyses (data not shown).

DISCUSSION

This is the most comprehensive and up to date systematic review and meta-analysis examining the impact of care at CACs compared with other hospitals on patient outcomes from OHCA. This review included data from large registries contributed by different countries. Patients cared for at CACs had increased likelihood of survival to hospital discharge and survival to hospital discharge with favourable neurological outcomes. However, there was no evidence that care at CACs improve survival to 30 days and survival to 30 days with favourable neurological outcomes. For patients with on-going CPR, care at CACs did not improve post-hospital arrival ROSC compared with other hospitals but this may be due to limited study size.

Whilst our findings are generally supportive of transporting OHCA patients to CACs, they should be interpreted with caution. Pooled data were extracted from retrospective and before-after studies, which all suffered from moderate or serious risk of bias. Study data was collected from different countries each with their distinctive healthcare systems and unique differences in system design, clinical practice, resuscitation and transporting policy of emergency medical services. Whilst some studies presented adjusted data and multivariate analyses, it may not be possible to adjust for unknown or unanticipated confounding factors, thereby limiting the internal validity of source evidence. The overall certainty of evidence was very low, preventing us from making firm conclusions

based on our findings. In addition, most studies conducted thus far have focused on care at a CAC per se, rather than on the more important question of whether patients that have an OHCA close to a non-CAC should be redirected for direct transportation to a CAC. Such studies conducted in different settings should include a more detailed analysis on possible problems associated with longer transportation times.

Relevant issues include the safety and logistics of longer prehospital transport time, the risk of re-arrest and other adverse events, the negative impact of transport on CPR quality, and whether to bypass a local hospital or arrange for secondary transfer after stabilization at a local hospital. [52-54] Our review did not specifically investigate the impact of transport duration on patient outcomes but did include some studies that reported transport times. Using a mathematical model, Kragholm, et al. concluded that there is survival benefit for cardiac arrest patients with prehospital ROSC to be transported to CAC even if journey time exceeds 20 minutes. [46] Similar findings were reported by Tranberg et al. who also reported that distances from arrest location to CAC was not associated with patient survival. [43] In contrast, regression analyses from Cournoyer study stated that increasing the delay from call to hospital arrival by 14.0 min would offset the potential survival benefit of being transported to a PCI-capable centre.[50] The decision whether to transport a patient and the hospital destination are likely to differ in urban, suburban, and rural settings. [55] Appropriately validated decision-making and risk prediction tools may help guide rescuers with transport and treatment decisions in future studies. [56, 57] Future clinical trials should also include the critical outcome of long term survival with favourable neurological outcomes to address this research gap.[7]

There was only one pilot randomised controlled study which reported only feasibility outcomes and did not demonstrate a difference between 30-day survival (CAC 9/18 50% vs other 6/15 40%) or 30-day survival with favourable neurological outcome (CAC 9/18 50% vs other 7/14 50%). [30] This study has just started recruiting. High quality evidence is needed to test the impact of CAC post cardiac arrest care on patient outcomes and address several remaining clinical knowledge gaps. For instance, specific subgroups of subjects (e.g. cardiac aetiology, shockable initial cardiac rhythm, ROSC or no ROSC prior to hospital admission) may benefit more or less from regional CAC care.

A key limitation is the general lack of consensus on what constitutes a CAC. In light of this, we accepted the study authors' definitions and descriptions of regional hospitals with a large case volume and coronary intervention capacity as a CAC. Likewise, we have not been able to substantiate or disprove these stated levels of service or other touted elements of post arrest care. We urge the Utstein writing group to consider CAC definition in the next iteration of Utstein variables for OHCA. Whilst clinical expertise and post arrest intervention should be readily available in CAC, hospital admission does not

necessarily mean that patients receive all elements of post arrest care. Patient care may be affected by other healthcare system pressures such as lack of resources or availability. Further considerations should also be given to hospital characteristics such as the number of cardiac arrest cases and other post-resuscitation hospital processes.[34, 58-61] A substantial challenge remains in identifying those components of post cardiac arrest care provided by a CAC that improve patient outcomes, especially recognizing the spectrum of disease severity across the post cardiac arrest syndrome and potential for tailored bundles of care.

From the included studies, the volume and annual case load of cardiac arrest did not improve patient survival in either CAC or other hospital setting. Mumma, et al. examined patient survival and neurological outcomes in CACs with different annual volume of cases (≤ 40 cases/year vs ≥ 40 cases/year) and found both survival and neurological outcomes of cardiac arrest patients were higher in CACs regardless of volume status.[51] There was also no reported difference in outcomes by hospital volume status (2 studies in CAC [40, 45], 1 study other hospitals [35], 1 study across all hospitals[38]).

Additional logistical and organization knowledge gaps remain in post cardiac arrest care. Regionalised post-cardiac arrest care is not feasible without a robust supporting network and regionalisation of medical services at the local and municipal levels.[62, 63] Fortunately, the trauma, STEMI, and stroke systems of care provide a template to establish a similar infrastructure for cardiac arrest. The success of whole system change such as those seen in major trauma systems cannot be ignored, and it needs to be highlighted that this change was not based on data from high quality randomized controlled trials but on observational data.[64] CAC services could potentially overlap with the cardiovascular and neurologic services offered at regional STEMI and stroke centres. Whilst indirect evidence established the cost effectiveness of stroke and STEMI centres, [65, 66] the cost effectiveness of specific strategies focusing on cardiac arrest patients and their potential impact on the care of other patient groups remains unknown.

The studies included in our review focused on clinical outcomes such as survival that has traditionally been considered to be of critical importance. Through the work of Core Outcome Set for Cardiac Arrest, there is a clear need for future research to address outcomes that are relevant and important to patient and their family such as quality of life and organ donation. [7, 67]

Limitations

We encountered high statistical heterogeneity in our meta-analysis and scrutinised included study characteristics for potential source of bias. Four studies [38, 40, 45, 46] included only patients with

prehospital ROSC in their analysis, possibly due to local policy and clinical practice.[68, 69] Survival from patients with no pre-hospital ROSC and requiring ongoing CPR may be better than commonly believed.[70] Before-and-after study designs used in four studies [36, 37, 48, 49] may not adequately control for changes in clinical practice. Both sets of sensitivity analyses did not adequately explain statistical heterogeneity. There was insufficient data from included studies for us to examine specifically whether there is any difference in outcomes for patients who present with shockable vs. non-shockable initial cardiac rhythm. We were also unable to conclusively assess the impact of direct transfer vs. secondary transfer to CAC. There was also no meaningful data to examine whether care at CAC has any influence on patient outcomes from in-hospital cardiac arrest compared to other hospitals.

CONCLUSIONS

Very low certainty evidence suggests that post-cardiac arrest care at cardiac arrest centres is associated with improved survival with favourable neurological outcome at hospital discharge and improved survival to hospital discharge. Care at CACs did not improve survival to 30 days with favourable neurological outcome and survival to 30 days. There remains a need of high quality data individual patient data meta-analysis and or data from randomised trials to fully elucidate the impact of CAC.

CONFLICTS OF INTEREST

Joyce Yeung was compensated by ILCOR for her work related to this review. TM and JB were authors of two studies included in this review. The other authors declare no competing conflict of interests.

ACKNOWLEDGMENTS

This Systematic Review was funded by the American Heart Association, on behalf of The International Liaison Committee on Resuscitation (ILCOR) for manuscript submission to the editor. The following authors received payment from this funding source to complete this systematic review:

Joyce Yeung as Expert Systematic Reviewer.

Nazi Torabi and Glyneva Bradley-Ridout as Information Services, St Michael's Hospital.

JY is supported by National Institute of Health Research Post-Doctoral Fellowship. JB is funded by a Heart Foundation Fellowship (#101171)

The authors would also like to acknowledge the contribution of Dr Eunice Singletary (ILCOR Continuous Evidence Evaluation liaison), Dr Robert Grief, Dr Farhan Bhanji (Education, Implementation and Team Task Force co-chairs) and Education, Implementation and Team Task Force and Advanced Life Support Task Force members.

CONFLICTS OF INTEREST

TM and JB were authors of two studies included in this review. Markus Skrifvars has received lecture fees from BARD MEDICAL and a research grant from GE Healthcare and INVOS COVIDIEN. The other authors declare no competing conflict of interests.

FUNDING ACKNOWLEDGMENT

This Systematic Review was funded by the American Heart Association, on behalf of The International Liaison Committee on Resuscitation (ILCOR). The following authors received payment from this funding source to complete this systematic review: Joyce Young as Expert Systematic Reviewer, Nazi Torabi and Glyneva Bradley-Ridout at Information Services, St Michael's Hospital.

ACKNOWLEDGEMENTS

JY is supported by National Institute of Health Research Post-Doctoral Fellowship. JB is funded by a Heart Foundation Fellowship (#101171).

The authors would also like to acknowledge the contribution of Dr Eunice Singletary (ILCOR Continuous Evidence Evaluation liaison), Dr Robert Grief, Dr Farhan Bhanji (Education, Implementation and Team Task Force co-chairs) and Education, Implementation and Team Task Force and Advanced Life Support Task Force members.

REFERENCES

- [1] Berdowski J, Berg RA, Tijssen JGP, Koster RW. Global incidences of out-of-hospital cardiac arrest and survival rates: Systematic review of 67 prospective studies. *Resuscitation*. 2010;81:1479-87.
- [2] Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. American heart association statistics committee and stroke statistics subcommittee. Heart disease and stroke statistics–2015 update: a report from the American Heart Association *Circulation*. 2015;131:e29-e322.
- [3] Gräsner J-T, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, et al. EuReCa ONE 27 Nations, ONE Europe, ONE Registry: A prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation*. 2016;105:188-95.
- [4] Beck B, Bray J, Cameron P, Smith K, Walker T, Grantham H, et al. Regional variation in the characteristics, incidence and outcomes of out-of-hospital cardiac arrest in Australia and New Zealand: Results from the Aus-ROC Epistery. *Resuscitation*. 2018;126:49-57.
- [5] Nichol G, Thomas E, Callaway CW, Hedges J, Powell JL, Aufderheide TP, et al. Regional Variation in Out-of-Hospital Cardiac Arrest Incidence and Outcome. *JAMA*. 2008;300:1423-31.
- [6] van Diepen S, Girotra S, Abella BS, Becker LB, Bobrow BJ, Chan PS, et al. Multistate 5-Year Initiative to Improve Care for Out-of-Hospital Cardiac Arrest: Primary Results From the HeartRescue Project. *Journal of the American Heart Association*. 2017;6:e005716.
- [7] Haywood K, Whitehead L, Nadkarni VM, Achana F, Beesems S, Böttiger BW, et al. COSCA (Core Outcome Set for Cardiac Arrest) in Adults: An Advisory Statement From the International Liaison Committee on Resuscitation. *Circulation*. 2018;CIR. 0000000000000562.
- [8] Nolan JP, Soar J. Post resuscitation care--time for a care bundle? *Resuscitation*. 2008;76:161-2.
- [9] Dumas F, White L, Stubbs BA, Cariou A, Rea TD. Long-term prognosis following resuscitation from out of hospital cardiac arrest: role of percutaneous coronary intervention and therapeutic hypothermia. *Journal of the American College of Cardiology*. 2012;60:21-7.
- [10] Søreide E, Busch M. Do cardiac arrest centres save more lives? Trends in Anaesthesia and Critical Care. 2016;7:21-5.
- [11] Sinha SS, Chen LM, Nallamothu BK. Survival by the Fittest: Hospital-Level Variation in Quality of Resuscitation Care. *Am Heart Assoc*; 2014. p. e000768.
- [12] Carr BG, Goyal M, Band RA, Gaieski DF, Abella BS, Merchant RM, et al. A national analysis of the relationship between hospital factors and post-cardiac arrest mortality. 2009;35:505-11.
- [13] Minei JP, Fabian TC, Guffey DM, Newgard CD, Bulger EM, Brasel KJ, et al. Increased trauma center volume is associated with improved survival after severe injury: results of a Resuscitation Outcomes Consortium study. *Annals of surgery*. 2014;260:456.
- [14] Tung Y-C, Chang G-M, Chien K-L, Tu Y-K. The relationships among physician and hospital volume, processes, and outcomes of care for acute myocardial infarction. *Medical care*. 2014;52:519-27.
- [15] Xian Y, Holloway RG, Chan PS, Noyes K, Shah MN, Ting HH, et al. Association between stroke center hospitalization for acute ischemic stroke and mortality. *Jama*. 2011;305:373-80.
- [16] MacKenzie EJ, Rivara FP, Jurkovich GJ, Nathens AB, Frey KP, Egleston BL, et al. A National Evaluation of the Effect of Trauma-Center Care on Mortality. *New England Journal of Medicine*. 2006;354:366-78.
- [17] Bhanji F, Finn JC, Lockey A, Monsieurs K, Frengley R, Iwami T, et al. Part 8: education, implementation, and teams: 2015 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation*. 2015;132:S242-S68.
- [18] Finn JC, Bhanji F, Lockey A, Monsieurs K, Frengley R, Iwami T, et al. Part 8: Education, implementation, and teams: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation*. 2015;95:e203-e24.
- [19] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339.

- [20] Tong JT, Eyngorn I, Mlynash M, Albers GW, Hirsch KG. Functional Neurologic Outcomes Change Over the First Six Months after Cardiac Arrest. *Critical care medicine*. 2016;44:e1202-e7.
- [21] Stiell IG, Nesbitt LP, Nichol G, Maloney J, Dreyer J, Beaudoin T, et al. Comparison of the Cerebral Performance Category score and the Health Utilities Index for survivors of cardiac arrest. *Annals of emergency medicine*. 2009;53:241-8. e1.
- [22] Nolan JP, Soar J, Cariou A, Cronberg T, Moulaert VRM, Deakin CD, et al. European Resuscitation Council and European Society of Intensive Care Medicine Guidelines for Post-resuscitation Care 2015. *Resuscitation*. 2015;95:202-22.
- [23] Higgins J, Green S. *Cochrane Handbook of Systematic Reviews of Interventions*. Version 5.1.0 [updated March 2011]. 2011.
- [24] Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *Bmj*. 2016;355:i4919.
- [25] Deeks JJ HJ, Altman DG. Chapter 9: Analysing Data and Undertaking Meta-Analyses. In: Higgins JPT GS, editor. *Cochrane Handbook for Systematic Reviews of Interventions Version 501: The Cochrane Collaboration*; 2008.
- [26] Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. 2003;327:557-60.
- [27] Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ, et al. What is "quality of evidence" and why is it important to clinicians? *BMJ*. 2008;336:995-8.
- [28] Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ (Clinical research ed)*. 2008;336:924-6.
- [29] Review Manager (RevMan) [Computer program]. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration. 2014.
- [30] Patterson T, Perkins GD, Joseph J, Wilson K, Van Dyck L, Robertson S, et al. A Randomised trial of Expedited transfer to a cardiac arrest centre for non-ST elevation ventricular fibrillation out-of-hospital cardiac arrest: the ARREST pilot randomised trial. *Resuscitation*. 2017;115:185-91.
- [31] Andrew E, Nehme Z, Wolfe R, Bernard S, Smith K. Long-term survival following out-of-hospital cardiac arrest. *Heart*. 2017;103:1104-10.
- [32] Elmer J, Rittenberger JC, Coppler PJ, Guyette FX, Doshi AA, Callaway CW, et al. Long-term survival benefit from treatment at a specialty center after cardiac arrest. *Resuscitation*. 2016;108:48-53.
- [33] Elmer J, Callaway CW, Chang C-CH, Madaras J, Martin-Gill C, Nawrocki P, et al. Long-Term Outcomes of Out-of-Hospital Cardiac Arrest Care at Regionalized Centers. *Annals of emergency medicine*. 2018;[In Press].
- [34] Brooks SC, Scales DC, Pinto R, Dainty KN, Racz EM, Gaudio M, et al. The Postcardiac Arrest Consult Team: Impact on Hospital Care Processes for Out-of-Hospital Cardiac Arrest Patients. *Critical Care Medicine*. 2016;44:2037-44.
- [35] Matsuyama T, Kiyohara K, Kitamura T, Nishiyama C, Nishiuchi T, Hayashi Y, et al. Hospital characteristics and favourable neurological outcome among patients with out-of-hospital cardiac arrest in Osaka, Japan. *Resuscitation*. 2017;110:146-53.
- [36] Tagami T, Hirata K, Takeshige T, Matsui J, Takinami M, Satake M, et al. Implementation of the fifth link of the chain of survival concept for out-of-hospital cardiac arrest. *Circulation*. 2012;126:589-97.
- [37] Seiner J, Polasek R, Lejsek J, Strycek M, Karasek J. Cardiac arrest center - One-year experience of the Regional Hospital Liberec. *Cor et Vasa*. 2018;60:e234-e8.
- [38] Couper K, Kimani PK, Gale CP, Quinn T, Squire IB, Marshall A, et al. Patient, health service factors and variation in mortality following resuscitated out-of-hospital cardiac arrest in acute coronary syndrome: Analysis of the Myocardial Ischaemia National Audit Project. *Resuscitation*. 2018;124:49-57.
- [39] Soholm H, Kjaergaard J, Bro-Jeppesen J, Hartvig-Thomsen J, Lippert F, Kober L, et al. Prognostic Implications of Level-of-Care at Tertiary Heart Centers Compared With Other Hospitals After Resuscitation From Out-of-Hospital Cardiac Arrest. *Circ Cardiovasc Qual Outcomes*. 2015;8:268-76.

- [40] Stub D, Smith K, Bray JE, Bernard S, Duffy SJ, Kaye DM. Hospital characteristics are associated with patient outcomes following out-of-hospital cardiac arrest. *Heart*. 2011;97:1489-94.
- [41] McKenzie N, Williams TA, Ho KM, Inoue M, Bailey P, Celenza A, et al. Direct transport to a PCI-capable hospital is associated with improved survival after adult out-of-hospital cardiac arrest of medical aetiology. *Resuscitation*. 2018;128:76-82.
- [42] Soholm H, Wachtell K, Nielsen SL, Bro-Jeppesen J, Pedersen F, Wanscher M, et al. Tertiary centres have improved survival compared to other hospitals in the Copenhagen area after out-of-hospital cardiac arrest. *Resuscitation*. 2013;84:162-7.
- [43] Tranberg T, Lippert FK, Christensen EF, Stengaard C, Hjort J, Lassen JF, et al. Distance to invasive heart centre, performance of acute coronary angiography, and angioplasty and associated outcome in out-of-hospital cardiac arrest: a nationwide study. *European Heart Journal*. 2017;38:1645-52.
- [44] Harnod D, Ma MHM, Chang WH, Chang RE, Chang CH. Mortality factors in out-of-hospital cardiac arrest patients: A nationwide population-based study in Taiwan. *International Journal of Gerontology*. 2013;7:216-20.
- [45] Chocron R, Bougouin W, Beganton F, Juvin P, Loeb T, Adnet F, et al. Are characteristics of hospitals associated with outcome after cardiac arrest? Insights from the Great Paris registry. *Resuscitation*. 2017;118:63-9.
- [46] Kragholm K, Malta Hansen C, Dupre ME, Xian Y, Strauss B, Tyson C, et al. Direct Transport to a Percutaneous Cardiac Intervention Center and Outcomes in Patients With Out-of-Hospital Cardiac Arrest. *Circ Cardiovasc Qual Outcomes*. 2017;10.
- [47] Lai CY, Lin FH, Chu H, Ku CH, Tsai SH, Chung CH, et al. Survival factors of hospitalized out-of-hospital cardiac arrest patients in Taiwan: A retrospective study. *PLoS ONE [Electronic Resource]*. 2018;13:e0191954.
- [48] Lick CJ, Aufderheide TP, Niskanen RA, Steinkamp JE, Davis SP, Nygaard SD, et al. Take Heart America: A comprehensive, community-wide, systems-based approach to the treatment of cardiac arrest. *Critical Care Medicine*. 2011;39:26-33.
- [49] Spaite DW, Bobrow BJ, Stolz U, Berg RA, Sanders AB, Kern KB, et al. Statewide regionalization of postarrest care for out-of-hospital cardiac arrest: association with survival and neurologic outcome. *Annals of Emergency Medicine*. 2014;64:496-506.e1.
- [50] Cournoyer A, Notebaert E, de Montigny L, Ross D, Cossette S, Londei-Leduc L, et al. Impact of the direct transfer to percutaneous coronary intervention-capable hospitals on survival to hospital discharge for patients with out-of-hospital cardiac arrest. *Resuscitation*. 2018;125:28-33.
- [51] Mumma BE, Diercks DB, Wilson MD, Holmes JF. Association between treatment at an ST-segment elevation myocardial infarction center and neurologic recovery after out-of-hospital cardiac arrest. *American Heart Journal*. 2015;170:516-23.
- [52] Olasveengen TM, Wik L, Steen PA. Quality of cardiopulmonary resuscitation before and during transport in out-of-hospital cardiac arrest. *Resuscitation*. 2008;76:185-90.
- [53] Kurz MC, Dante SA, Puckett BJ. Estimating the impact of off-balancing forces upon cardiopulmonary resuscitation during ambulance transport. *Resuscitation*. 2012;83:1085-9.
- [54] Becker LR, Zaloshnja E, Levick N, Li G, Miller TR. Relative risk of injury and death in ambulances and other emergency vehicles. *Accident Analysis & Prevention*. 2003;35:941-8.
- [55] Geri G, Gilgan J, Wu W, Vijendira S, Ziegler C, Drennan IR, et al. Does transport time of out-of-hospital cardiac arrest patients matter? A systematic review and meta-analysis. *Resuscitation*. 2017;115:96-101.
- [56] Poppe M, Weiser C, Holzer M, Sulzgruber P, Datler P, Keferböck M, et al. The incidence of "load&go" out-of-hospital cardiac arrest candidates for emergency department utilization of emergency extracorporeal life support: A one-year review. *Resuscitation*. 2015;91:131-6.
- [57] Drennan IR, Lin S, Sidalak DE, Morrison LJ. Survival rates in out-of-hospital cardiac arrest patients transported without prehospital return of spontaneous circulation: an observational cohort study. *Resuscitation*. 2014;85:1488-93.

- [58] Schober A, Sterz F, Laggner AN, Poppe M, Sulzgruber P, Lobmeyr E, et al. Admission of out-of-hospital cardiac arrest victims to a high volume cardiac arrest center is linked to improved outcome. 2016;106:42-8.
- [59] Callaway CW, Schmicker R, Kampmeyer M, Powell J, Rea TD, Daya MR, et al. Receiving hospital characteristics associated with survival after out-of-hospital cardiac arrest. Resuscitation. 2010;81:524-9.
- [60] Callaway CW, Schmicker RH, Brown SP, Albrich JM, Andrusiek DL, Aufderheide TP, et al. Early coronary angiography and induced hypothermia are associated with survival and functional recovery after out-of-hospital cardiac arrest. Resuscitation. 2014;85:657-63.
- [61] Stub D, Schmicker RH, Anderson ML, Callaway CW, Daya MR, Sayre MR, et al. Association between hospital post-resuscitative performance and clinical outcomes after out-of-hospital cardiac arrest. Resuscitation. 2015;92:45-52.
- [62] Donnino MW, Rittenberger JC, Gaieski D, Cocchi MN, Giberson B, Peberdy MA, et al. The development and implementation of cardiac arrest centers. Resuscitation. 2011;82:974-8.
- [63] Martin-Gill C, Dilger CP, Guyette FX, Rittenberger JC, Callaway CW. Regional impact of cardiac arrest center criteria on out-of-hospital transportation practices. Prehospital emergency care : official journal of the National Association of EMS Physicians and the National Association of State EMS Directors. 2011;15:381-7.
- [64] Moran CG, Lecky F, Bouamra O, Lawrence T, Edwards A, Woodford M, et al. Changing the System-Major Trauma Patients and Their Outcomes in the NHS (England) 2008–17. 2018.
- [65] Guzauskas GF, Boudreau DM, Villa KF, Levine SR, Veenstra DLJS. The cost-effectiveness of primary stroke centers for acute stroke care. 2012;43:1617-23.
- [66] Regueiro A, Bosch J, Martin-Yuste V, Rosas A, Faixedas MT, Gomez-Hospital JA, et al. Cost-effectiveness of a European ST-segment elevation myocardial infarction network: results from the Catalan Codi Infart network. BMJ Open. 2015;5:e009148.
- [67] Haywood KL, Pearson N, Morrison LJ, Castren M, Lilja G, Perkins GD. Assessing health-related quality of life (HRQoL) in survivors of out-of-hospital cardiac arrest: A systematic review of patient-reported outcome measures. Resuscitation. 2018;123:22-37.
- [68] Kellermann AL, Staves DR, Hackman BB. In-hospital resuscitation following unsuccessful prehospital advanced cardiac life support: 'heroic efforts' or an exercise in futility? Annals of emergency medicine. 1988;17:589-94.
- [69] Kellerman A, Hackman B, Somes G. Cardiology: Survival Poor After Unsuccessful Prehospital Advanced Cardiac Life Support. Physician Assistant. 1993;17:13-9.
- [70] Eisenburger P, Havel C, Sterz F, Uray T, Zeiner A, Haugk M, et al. Transport with ongoing cardiopulmonary resuscitation may not be futile. Br J Anaesth. 2008;101:518-22.

LEGENDS TO FIGURES

Figure 1 PRISMA Diagram




Figure 2 Survival to 30 days with favourable outcome. Higher Odds Ratio favours CAC.




Figure 3 Survival to hospital discharge with favourable outcome. Higher Odds Ratio favours CAC.




Figure 4 Survival to 30 days. Higher Odds Ratio favours CAC.




Figure 5 Survival to hospital discharge. Higher Odds Ratio favours CAC.




Figure 6 Return of spontaneous circulation post-hospital arrival in patients with ongoing resuscitation. Higher Odds Ratio favours CAC.




Table 1 Characteristics of included studies

| Study | Setting | Study Design | Sample size | Population | Description of CAC | Key relevant findings |
|----------------|--------------------------------|--|---|--|--|---|
| Andrew 2017** | Victoria, Australia | Retrospective cohort using registry data 2000-2014 Victoria Ambulance Cardiac Arrest registry | Total 3449 CAC 2775 nonCAC 674 | All OHCA survivors to hospital discharge | PCI capable hospital | <ul style="list-style-type: none"> Care at CAC was associated with long term survival post hospital discharge in multivariate analyses |
| Brooks 2016** | Canada | Prospective cohort study 2011-2013 | Total 1006 CAC (2 hospitals with Post Arrest Consult Team, PACT) 151 nonCAC (no PACT) 855 | Non traumatic OHCA, aged >18, GCS<10 on ED arrival, survive for 6hrs | Post arrest consult team (PACT) On call doctor and nurse 24/7, review every 24hrs Advice on TTM, PCI, electrophysiology assessment, neuroprognostication | <ul style="list-style-type: none"> PACT hospital was not associated with survival to hospital discharge or survival with good neurologic status (mRS 0-2) in adjusted analyses |
| Chocron 2017 | Paris, France | Retrospective analysis from database Paris Sudden Death Expertise Centre 2011-2013 | Total 1436 CAC 917 nonCAC 519 | Non traumatic >18, OHCA achieved ROSC, admitted alive | High case volume with PCI 24/7 | <ul style="list-style-type: none"> Care at CAC was associated with survival to discharge in univariate analysis but not in multivariate analyses |
| Couper 2018 | United Kingdom | Retrospective analysis using Myocardial Ischaemia National Audit Project Database | Total 17604 PCI centres 7800 nonPCI centres 9804 | Aged >18, OHCA, achieved ROSC prehospital care | PCI centres performing >100 cases/year | <ul style="list-style-type: none"> Care at CAC was associated with survival to discharge in univariate analysis but not in multivariate analyses |
| Cournoyer 2018 | Montreal, Canada | Retrospective cohort study using a registry of OHCA from the region of Montreal, Canada April 2010 until December 2015 | Total 4922 CAC 2389 nonCAC 2533 | Aged >18, All transported non-traumatic OHCA without 'do-not-resuscitate' directives or with 'obviously deceased' criteria | PCI-capable hospital (STEMI centre) with PCI or hemodynamic support 24/7 | <ul style="list-style-type: none"> Care at CAC was associated with survival to discharge in adjusted analyses |
| Elmer 2016** | Southwestern Pennsylvania, USA | Retrospective analysis from database 2005-2013 IHCA and OHCA | Total 987 CAC (1 CAC) 680 nonCAC (2 moderate volume tertiary centres 4 low volume centres) 307 | Aged>18, IHCA and OHCA who survived to discharge (exclude trauma, neurological catastrophe) | Regional referral centre organised post arrest care | <ul style="list-style-type: none"> Care at high volume CAC was associated with survival to hospital discharge with good neurological outcome in adjusted analyses |
| Elmer 2018# | Southwestern Pennsylvania, USA | Retrospective analysis from EMS database 2010-2014 OHCA | Total 5217 CAC 920 nonCAC 4297 <ul style="list-style-type: none"> Direct 570 Secondary 350 | Aged>16yrs, OHCA with prehospital ROSC, including secondary transfer | 'Cardiac arrest receiving centre' – receiving at least 1 interfacility transfer OHCA patient every 3 months | <ul style="list-style-type: none"> Care at CAC was associated with long term survival in adjusted analyses |

| | | | | | | |
|-----------------------------|---|--|--|--|--|--|
| Harnod 2013 | Taiwan | Retrospective cohort using National Health Insurance Research database 2005-2007 | Total 1673 CAC (22 medical centres) 435 nonCAC (72 Regional hospitals, 367 district hospitals) 1238 | Aged >15yrs, OHCA (exclude Trauma, without intubation or CPR in ED) | 'Medical centres' | <ul style="list-style-type: none"> Care at CAC was associated with survival at 30 days after hospital discharge in adjusted analyses |
| Kragholm 2017 | USA Cardiac Arrest Registry to Enhance Survival (CARES) | Retrospective analysis from CARES database 2012-2014 | Total 3449 CAC 1359 nonCAC 148 | Arrests of presumed cardiac cause with prehospital ROSC | PCI centre (primary PCI was available on a 24/7 basis) | <ul style="list-style-type: none"> Care at CAC was associated with improved survival to hospital discharge with good neurological outcome and survival to hospital discharge in adjusted analyses |
| Lai 2018 | Taiwan | Retrospective cohort using National Health Insurance Research database 2007 to 2013 | Total 11000 CAC 2255 nonCAC 8745 | OHCA with codes of VF and cardiac arrest (exclude <18, Unknown outcomes, trauma) | Medical centres | <ul style="list-style-type: none"> Care at CAC is associated with survival to hospital discharge in adjusted analyses |
| Lick 2011 | Minnesota, USA | Before and after (2005 compared with 2006-2007) | Total 353 CAC 247 nonCAC 106 | All OHCA | Take Heart Program Cardiac arrest centres/Resuscitation centre of excellence: TTM, 24/7 PCI, electrophysiology, outcome tracking | <ul style="list-style-type: none"> Care at CAC was associated with survival to hospital discharge in adjusted analyses Care at CAC was not associated with improved mean CPC score at discharge in adjusted analyses |
| Matsuyama 2017 [#] | Osaka, Japan | Retrospective analysis Utstein Osaka Project 2005-2012 | Total 44,474 CAC (16CCMC) 17,737 nonCAC (301 non-CCMC) 26,737 | Aged >18, OHCA, resuscitated by EMS and brought to hospital, all causes | Critical care medical centre: ≥20 beds and ICU for critically ill patients, capable of ECPR or PCI and TTM 24/7. | <ul style="list-style-type: none"> Care at CAC was associated with survival at 30 days with good neurological outcome, survival at 30 days and ROSC in adjusted analyses |
| McKenzie 2018 | Perth, Australia | Retrospective analysis St John Ambulance Western Australia OHCA Database January 2012 to December 2015 | Total 539 Non CAC 26 CAC 513 <ul style="list-style-type: none"> Direct 408 Secondary 105 | Aged >18, OHCA, admitted to and survived ED care | 24/7 PCI centre and post resuscitation care | <ul style="list-style-type: none"> Care at CAC was associated with survival to hospital discharge in adjusted analyses Direct transport to CAC was associated with survival to hospital discharge in adjusted analyses Indirect transport to CAC was associated with increased risk of death up to 12-months in adjusted analyses |
| Mumma 2015 | California, USA | Retrospective cohort using registry data 2011 | Total 7725 CAC (125 STEMI centre) 5202 nonCAC (208 non-STEMI centre) 2523 | Discharge database with cardiac arrest on care | STEMI centre: 24/7 PCI and TTM, >40 patients/yr ROSC post OHCA | <ul style="list-style-type: none"> Care at CAC was associated with survival at hospital discharge with good neurological outcome (defined as discharge to home, residential care facility, prison, jail, another hospital for |

| | | | | | | |
|-----------------|---------------------|--|---|---|--|---|
| | | | | | | nonacute care, left against advice) in adjusted analyses |
| Patterson 2017* | London, UK | Pilot study of Randomised controlled trial | Total 33 CAC (one hospital) 18 nonCAC (6 hospitals) 15 | Aged >18, OHCA , witnessed VF presumed cardiac cause, attended by advanced paramedic practitioners | 24/7 PCI,GDT, TTM | <ul style="list-style-type: none"> Care at CAC was not associated with survival at 30 days with good neurological outcome, survival at 30 days in adjusted analyses (pilot outcome only) |
| Seiner 2018 | Liberec, Czech | Before and after study 2013-2015 compared to 2016-2017 | Total 33 CAC (After) 61 nonCAC (Before) 47 | Aged >18, OHCA , all cause | Cardiovascular centre with 24/7 cath lab, PCAS | <ul style="list-style-type: none"> Care at CAC was not associated with survival at 30 days with good neurological outcome and survival at 30 days in unadjusted analyses |
| Soholm 2013 | Copenhagen, Denmark | Retrospective registry 2002-2010 | Total 1020 CAC 563 nonCAC 457 | OHCA with or without ROSC, dispatch of paramedic & anaesthetist include non-cardiac Excludes STEMI | Tertiary hospital 24/7 cath lab, cardiac ITU, TTM (from 2004) | <ul style="list-style-type: none"> Care at CAC was associated with survival to 30 days in adjusted analyses |
| Soholm 2015 | Copenhagen, Denmark | Retrospective analysis 2002-2011 | Total 33 CAC (2 tertiary university heart centres) 586 nonCAC (6 non-tertiary university hospital) 492 | Aged >18, OHCA transported with ROSC or ongoing CPR Excludes STEMI | Tertiary centre: heart centre with dedicated anaesthetists, cardiologists and surgeons. Interventional cardiology and cardiac surgery 24/7, cardiac ICU, TTM (2002-2004) | <ul style="list-style-type: none"> Care at CAC was associated with survival to hospital discharge and survival at 30 days in adjusted analyses Care at CAC was not associated with survival to hospital discharge with good neurological outcome in adjusted analyses |
| Spaite 2014 | Arizona, USA | Before and after 2007-2010 After: (December 14, 2007, to November 25, 2010) Before: (January 1, 2007, and December 13, 2007) | Total 2177 CAC 1737 nonCAC 440 | Aged >18, OHCA presumed cardiac transported | Cardiac receiving centre Coronary angiography/PCI, TTM, Statewide regionalisation | <ul style="list-style-type: none"> Care at CAC was associated with survival to hospital discharge and survival to hospital discharge with favourable neurological outcomes in adjusted analyses |
| Stub 2011 | Victoria, Australia | Retrospective analysis 2003-2010 Victoria ambulance data | Total 2706 CAC 1816 nonCAC 890 | Aged>18, OHCA presumed cardiac transported with ROSC | 24hr cardiac interventional service | <ul style="list-style-type: none"> Care at CAC was associated with survival to hospital discharge in adjusted analyses Care at CAC was associated with survival to hospital discharge with favourable neurological outcome (defined as discharge home) in unadjusted analyses |

| | | | | | | |
|----------------------------|-------------|---|-------------------------------------|---|---|---|
| Tagami 2012 | Aizu, Japan | Before and after 2006-2008 compared with 2009-2010 | Total 1482 CAC 712 nonCAC 770 | OHCA transported with ROSC or ongoing CPR | Post resuscitation care centre: Tertiary centre PCAS, TTM, PCI | <ul style="list-style-type: none"> Care at CAC was associated with survival at 30 days with good neurological outcome and survival to discharge in adjusted analyses |
| Tranberg 2017 [#] | Denmark | Retrospective analysis 2001-2013 Danish Cardiac Arrest Registry | CAC 900 nonCAC 1300 | All OHCA | High-volume invasive heart centres with a 24-h PCI service | <ul style="list-style-type: none"> Care at CAC was associated with survival at 30 days in adjusted analyses |

*Study data not included in meta-analysis; [#] Authors contacted to provide additional data

CAC: Cardiac arrest centre; nonCAC: other institution not designated as CAC; CPC: cerebral performance category; ECPR: extracorporeal CPR; ED: emergency department; GCS: Glasgow Coma Score; ICU: intensive care unit; IHCA: in-hospital cardiac arrest; OHCA: out-of-hospital cardiac arrest; PCAS: Post cardiac arrest service; PCI: Primary Coronary Intervention; QoL: Quality of life; ROSC: Return of spontaneous circulation STEMI: ST-elevation myocardial infarction; TTM: Targeted temperature management; VF: ventricular fibrillation

Table 2a Risk of bias – randomised study

| Study | Randomisation (selection bias) | Allocation concealment (selection bias) | Performance and Detection bias | Attrition bias | Reporting bias | Other bias |
|----------------|-----------------------------------|---|-----------------------------------|----------------|----------------|-------------|
| Patterson 2017 | LOW | HIGH | UNCLEAR | LOW | LOW | PILOT STUDY |

Table 2b Risk of bias – non randomised studies

| Study ID | Bias in confounding | Bias in patient selection | Bias in classification of intervention | Bias in deviation from intervention | Bias from missing data | Bias from measuring outcomes | Bias from selected reporting of results | Overall |
|----------------|------------------------|------------------------------|--|---|---------------------------|------------------------------------|--|----------------------|
| Andrew 2017 | Serious | Moderate | Low | Low | Moderate | Moderate | Low | Serious ^a |
| Brooks 2016 | Moderate | Low | Low | Low | Low | Moderate | Low | Moderate |
| Chocron 2017 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Couper 2018 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Cournoyer 2018 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Elmer 2016 | Serious | Serious | Low | Moderate | Low | Moderate | Low | Serious ^b |
| Elmer 2018 | Serious | Serious | Low | Low | Moderate | Moderate | Low | Serious ^b |
| Harnod 2013 | Serious | Moderate | Low | Low | Low | Moderate | Low | Serious ^c |
| Kragholm 2017 | Serious | Moderate | Low | Low | Moderate | Moderate | Low | Serious ^d |
| Lai 2018 | Serious | Serious | Low | Low | Low | Serious | Low | Serious ^e |
| Lick 2011 | Serious | Serious | Low | Low | Low | Serious | Low | Serious ^f |
| Matsuyama 2017 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Mckenzie 2018 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Mumma 2015 | Moderate | Moderate | Low | Low | Moderate | Moderate | Low | Moderate |
| Seiner 2018 | Serious | Low | Low | Low | Low | Moderate | Low | Serious ^g |
| Soholm 2013 | Serious | Moderate | Low | Low | Moderate | Moderate | Low | Serious ^h |
| Soholm 2015 | Serious | Serious | Low | Low | Low | Moderate | Moderate | Serious ⁱ |
| Spaite 2014 | Serious | Moderate | Low | Low | Moderate | Moderate | Low | Serious ^j |
| Stub 2011 | Moderate | Moderate | Low | Low | Moderate | Moderate | Low | Moderate |
| Tagami 2012 | Moderate | Moderate | Low | Low | Low | Moderate | Low | Moderate |
| Tranberg 2017 | Serious | Moderate | Low | Low | Moderate | Moderate | Low | Serious ^k |

^a only survivors were included^b only survivors were included, included interhospital transfer patients

^c inadequate adjustment of confounding factors

^d Difference in EMS response time, transfer times, whether bypass hospital, urban/rural arrest location

^e only those with prehospital ROSC included which is dependent on transfer time and other confounders; coding does not differentiate between pre-existing conditions and secondary diagnoses

^f a system of change including CPR training, bystander CPR, TTM, introduction of mechanical CPR device; differences in rate of patient recruitment before and after phase

^g inadequate adjustment for confounders

^h excluded all STEMI patients

ⁱ excluded all STEMI patients, included patients from 2013 paper

^j included CPR training program, EMS transfer rates

^k assumption of location of arrest in 23% of patients, significant temporal changes in bystander CPR rates